Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information. including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS. 1. REPORT DATE (DD-MM-YYYY) 2. REPORT TYPE 3. DATES COVERED (From - To) Technical Paper 4. TITLE AND SUBTITLE 5a. CONTRACT NUMBER 5b. GRANT NUMBER 5c. PROGRAM ELEMENT NUMBER 6. AUTHOR(S) 5d. PROJECT NUMBER 2303 5e. TASK NUMBER 5f. WORK UNIT NUMBER 345709 8. PERFORMING ORGANIZATION 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) REPORT 9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) 10. SPONSOR/MONITOR'S ACRONYM(S) Air Force Research Laboratory (AFMC) 11. SPONSOR/MONITOR'S AFRL/PRS NUMBER(S) 5 Pollux Drive Edwards AFB CA 93524-7048 12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited. 13. SUPPLEMENTARY NOTES 14. ABSTRACT 20030129 225 15. SUBJECT TERMS 16. SECURITY CLASSIFICATION OF: 18. NUMBER 19a. NAME OF RESPONSIBLE 17. LIMITATION OF ABSTRACT OF PAGES PERSON Leilani Richardson a. REPORT b. ABSTRACT c. THIS PAGE 19b. TELEPHONE NUMBER (include area code) (661) 275-5015 Unclassified Unclassified Unclassified Standard Form 298 (Rev. 8-98)

REPORT DOCUMENTATION PAGE

Form Approved

OMB No. 0704-0188

Prescribed by ANSI Std. 239.18

BORMERCE

MEMORANDUM FOR PRS (In-House/Contractor Publication)

FROM: PROI (STINFO)

01 Mar 2001

SUBJECT: Authorization for Release of Technical Information, Control Number: AFRL-PR-ED-TP-2001-042 Vij, Ashwani (ERC); Wilson, W.I, et al., "Polynitrogen Chemistry, Synthesis, Characterization, and Crystal Structure of Surprisingly Stable Fluorantimonate Salts of N₅⁺"

Journal of American Chemical Society (Deadline: N/A)

(Statement A)

Polynitrogen Chemistry. Synthesis, Characterization, and Crystal Structure of Surprisingly Stable Fluoroantimonate Salts of N_5^+

Ashwani Vij, † William W. Wilson, † Vandana Vij, † Fook S. Tham, § Jeffrey A. Sheehy, †
and Karl O. Christe, *†‡

on centering

Propulsion Sciences and Advanced Concepts Division, Air Force Research Laboratory (AFRL/PRS), Edwards AFB, California 93524, Loker Hydrocarbon Research Institute and Department of Chemistry, University of Southern California, Los Angeles, CA 90089, and Department of Chemistry, University of California, Riverside, CA 92521

Abstract

The new N_5^+ salt, N_5^+ SbF₆, was prepared from N_2F^+ SbF₆ and HN₃ in anhydrous HF solution. The white solid is surprisingly stable, decomposing only at 70 °C, and is relatively insensitive to impact. Its vibrational spectrum exhibits all nine fundamentals with frequencies that are in excellent agreement with the theoretical calculations for a five-atomic V-shaped ion of C_{2v} symmetry. The N_5^+ Sb₂F₁₁ salt was also prepared and its crystal structure was determined. The geometry previously predicted for free gaseous N_5^+ from theoretical calculations was confirmed within experimental error. The Sb₂F₁₁ anions exhibit an unusual geometry with eclipsed SbF₄ groups due to inter-ionic bridging with the N_5^+ cations. The N_5^+ cation is a powerful one-electron oxidizer. Its electron affinity falls between 11.0 and 12.08 eV because it readily oxidizes NO to NO⁺ and NO₂ to NO₂⁺ but fails to oxidize Xe or O₂.

Introduction

The recent discovery of $N_5^+AsF_6^-$ as a marginally stable compound that can be prepared on a macroscopic scale is quite remarkable. It represented only the third readily accessible homoleptic polynitrogen compound besides N_2 and N_3^- and as such has received much public acclaim. Since N_5AsF_6 is only marginally stable and had given rise to some explosions, it was of great interest to search for more stable N_5^+ salts in order to allow a more thorough characterization of this fascinating cation and to provide a suitable starting material for the pursuit of nitrogen allotropes. In this paper, the synthesis and characterization of surprisingly stable fluoroantimonate salts of N_5^+ and the crystal structure of $N_5^+Sb_2F_{11}^-$ are reported.

Experimental

Caution! HN_3 , azides and polynitrogen compounds are highly endothermic and can decompose explosively. They should be handled only on a small scale with appropriate safety precautions (face shields, leather gloves, and protective clothing). Condensation of neat HN_3 at -196 °C into Teflon ampoules containing oxidizers has repeatedly resulted in explosions upon condensation or melting of the HN_3 .

Materials and Apparatus. All reactions were carried out in ¾ inch o. d. Teflon-FEP or -PFA ampoules that contained Teflon coated magnetic stirring bars and were closed by stainless steel valves. Volatile materials were handled on a stainless steel / Teflon-FEP vacuum line. Nonvolatile solids were handled in the dry nitrogen atmosphere of a glove box. HN₃ was generated and handled on a Pyrex glass vacuum line

equipped with grease-free Kontes glass-Teflon valves. Infrared spectra were recorded on a Mattson Galaxy FT-IR spectrometer using dry powders pressed between AgCl windows in an Econo press (Barnes Engineering Co.). Raman spectra were recorded on a Bruker Equinox 55 FT-RA spectrometer using a Nd-Yag laser at 1064 nm and Pyrex melting point capillaries as sample containers. The thermal stabilities were determined using a DuPont Model 910 DSC, crimp-sealed aluminum pans as sample containers, and heating rates of 3 °C/min. The data were recorded and analyzed with a DuPont Model 2000 thermal analyst. Impact sensitivities were measured on an Olin Mathieson drop weight tester standardized with RDX (hexahydro-1,3,5-trinitro-1,3,5-triazine, 30 kg·cm, 50 %).

The N₂F⁺SbF₆ and N₂F⁺Sb₂F₁₁ starting materials were prepared from *cis*-N₂F₂ and SbF₅ in anhydrous HF solution as previously described.⁴⁻⁷ The HF (Matheson Co.) was dried by storage over BiF₅ (Ozark Mahoning).⁸ The NO and NO₂ (Matheson Co) were purified by fractional condensation prior to their use. The O₂ and Xe (Matheson Co) were used as received. The preparation of HN₃ has previously been described.¹

Preparation of N₅+SbF₆. A Teflon ampoule, equipped with a stainless steel valve and containing a Teflon-coated magnetic stirring bar, was passivated with ClF₃. It was attached to the metal vacuum line and treated several times with anhydrous HF until no color was observed upon freezing the HF at -196 °C. It was then loaded with N₂F⁺SbF₆ (4.97 mmol) in the glove box and attached to the metal vacuum line. The ampoule was evacuated and cooled to -196 °C. Anhydrous HF (~2 mL) was then condensed into the ampoule and its contents were allowed to warm to ambient temperature with occasional stirring. After all the N₂F⁺SbF₆ had dissolved, the ampoule

was re-cooled to -196 °C and some additional neat HF was condensed onto the upper walls of the tube where the HN₃ was going to be frozen out. The cold ampoule was then connected to the glass line, and HN₃ (5.00 mmol) was added slowly at -196 °C. The reaction mixture was allowed to warm slowly behind a safety shield to room temperature and kept at this temperature for about 45 min. The volatile materials were removed by pumping for several hours at 20 °C, leaving behind a white powder (1.502 g, weight calcd for 4.97 mmol of N₅SbF₆ = 1.520 g) that was identified by its vibrational spectra as N₅SbF₆.

This reaction was also carried out by first condensing HN_3 at -196 °C into a prepassivated and preweighed Teflon ampoule containing a known amount of HF. The resulting mixture was homogenized at ambient temperature. The ampoule was taken into the glove box, where a stoichiometric amount of $N_2F^+SbF_6^-$ was added at -196 °C. The cold ampoule was attached to the metal vacuum line and evacuated. Subsequent slow warming of the reaction mixture to room temperature for about 30 min, followed by removal of all volatile material resulted in the isolation of $N_5^+SbF_6^-$ in >99% yield.

Preparation of N₅+Sb₂F₁₁. Freshly distilled SbF₅ (1.449 mmol) was added in the glove box to a prepassivated Teflon-FEP ampoule, and HF (1.9 mL liquid) was added on the metal vacuum line at -196 °C. The mixture was homogenized at room temperature and taken back into the glove box. The ampoule was cooled inside the glove box to -196 °C and opened, and N₅+SbF₆ (1.444 mmol) was added. The resulting mixture was allowed to warm to room temperature, and all volatile material was pumped off. The white solid residue (758 mg, weight calcd. for 1.444 mmol of N₅+Sb₂F₁₁ = 755 mg) was shown by vibrational spectroscopy to consist of N₅+Sb₂F₁₁.

Reactions of $N_5^+SbF_6^-$ with NO, NO₂, O₂ or Xe. In a typical experiment, a .5 inch Teflon-FEP ampoule, that was closed by a Teflon valve, was loaded in the dry-box with $N_5^+SbF_6^-$ (.53 mmol). On the vacuum line, NO (4.2 mmol) was added at $-196\,^{\circ}$ C and the contents of the ampoule were allowed to warm slowly with intermittent cooling to room temperature. After keeping the ampoule for 2 hr at room temperature, it was cooled back to $-196\,^{\circ}$ C and the volatile gas (1.34 mmol of N_2) was measured and pumped off. The amount of unreacted NO was measured (3.6 mmol) and pumped off at room temperature, leaving behind .53 mmol of NO⁺SbF₆⁻ that was identified by vibrational spectroscopy.

In a similar manner, N_5 ⁺Sb F_6 ⁻ was found to react quantitatively with NO₂, but no reaction was observed with either Xe or O₂.

Crystal structure determination of N₅+Sb₂F₁₁. About 1 mL of anhydrous SO₂ was condensed onto 0.200 g of N₅SbF₆ at -196 °C in a ½ inch o. d. sapphire tube (Tyco Corp.) closed by a stainless steel valve. The contents of the tube were warmed to -78 °C causing all of the N₅SbF₆ to dissolve and form a pale yellowish solution. Anhydrous SO₂ClF (~1.5 mL) was then slowly condensed onto this solution under vacuum. The solvents were then slowly removed under a static vacuum at -64 °C over a period of ~16 hours leaving behind plate-like colorless crystals. These crystals were extremely reactive to perfluoropolyether oil and showed an instantaneous evolution of nitrogen gas. The majority of the crystals were very soft and difficult to handle, but a few crystals appeared to exhibit a different habit and better mechanical strength. One of these crystals was immersed in halocarbon grease and mounted on the goniometer head using a precentered Nylon Cryoloop equipped with a magnetic base. The structure of the salt was determined

using a Bruker diffractometer equipped with a CCD detector and a low temperature, LT3, device. The 3-circle platform with a fixed χ-axis was controlled by the SMART⁹ software package. The unit cell parameters were determined at -60 °C from three runs of data with 30 frames per run using a scan speed of 30-seconds per frame. A complete hemisphere of data was collected, using 1271 frames at 30 sec/frame, including 50 frames that were collected at the beginning and end of the data collection to monitor crystal decay. Data were integrated using the SAINT¹⁰ software package, and the raw data was corrected for absorption using the SADABS¹¹ program. The absence of h + k =odd and h0l reflections (l = odd) shows the presence of a C-centered lattice and a c-glide plane parallel and perpendicular to the b-axis, respectively, indicating Cc or C2/c as the likely space groups. The intensity statistics, E²-1 values, indicated a centrosymmetric space group thereby excluding Cc as a possible space group. The space group was thus unambiguously assigned as C2/c. The structure was solved by the Patterson method using the SHELXS-97¹² program and refined by the least squares method on F² using SHELXL-97.¹³ The initial Patterson map revealed the position of the two Sb atoms linked by a fluorine atom. The remaining atoms were located from subsequent difference electron density maps and finally refined anisotropically by the least-squares method on F² using the SHELXTL¹⁴ 5.1 software for Windows NT. The crystal did not show any significant decomposition during the data collection. The experimental and refinement parameters, and the atomic coordinates and thermal displacement parameters are listed in Tables 1 and 2, respectively.

Results and Discussion

 $N_2F^+SbF_6^-$

of $H_2N_3^+SbF_6^{-15}$

Synthesis and Properties of N₅+SbF₆. The synthesis of N₅+SbF₆ was carried out in the same manner as previously reported for N₅+AsF₆ by reacting N₂F⁺SbF₆ with HN₃ in anhydrous HF solution at -78 °C, followed by removal of the volatile products at room temperature.

$$N_2F^+SbF_6^-$$
 + HN_3 $N_5^+SbF_6^-$ + HF
The yield of $N_5^+SbF_6^-$ is essentially quantitative and the product purity is high. It is essential that the reaction system is completely anhydrous as water hydrolyzes the N_5^+ salt generating free SbF_5 which in combination with HF protonates HN_3 under formation

HF

 HN_3

The N₅⁺SbF₆⁻ salt is a colorless hygroscopic solid that is stable at ambient temperature and, based on the DSC data, starts to decompose at 70 °C. It is surprisingly insensitive to impact. Even at the maximum setting of our apparatus (200 kg·cm), only partial thermal decomposition due to adiabatic heating of the sample but no explosions were observed. The salt is soluble in and compatible with HF, SO₂, and CHF₃.

The oxidizing properties of N₅⁺SbF₆ were examined in the solid state and in HF solution by exposing it to 2 atmospheres of either oxygen or xenon gas between -78 °C and ambient temperature. No oxidations to O_2^+ and Xe_2^+ , respectively, were observed under these conditions, showing that the electron affinity of N₅⁺ is lower than the first ionization potential of xenon (12.08 eV), i. e., N₅⁺ is a weaker oxidizer than either PtF₆ that can oxidize O_2 to O_2^+ 16 or O_2^+ that can oxidize Xe to Xe_2^+ under similar conditions. 17,18 In spite of the first ionization potential of N₂ (15.51 eV) being 3.01 eV higher than that of O_2 (12.5 eV), the electron affinity of N_5^+ is lower than that of O_2^+

because in N_5^+ the positive charge is spread over a larger number of atoms, thereby decreasing its oxidizing power. However, it was found that N_5^+ can quantitatively oxidize either NO (1. IP = 9.5 eV) or NO₂ (1. IP = 11.0 eV) according to:

NO +
$$N_5^+SbF_6^-$$
 NO $^+SbF_6^-$ + 2.5 N_2
and NO₂ + $N_5^+SbF_6^-$ NO₂ $^+SbF_6^-$ + 2.5 N_2

Therefore the electron affinity of N_5^+ must have a value between 11.0 and 12.08 eV, rendering it a very strong one-electron oxidizer. Although it is not quite as powerful as PtF_6 or O_2^+ salts, it offers the great advantage of not acting as a fluorinating or oxygenating agent, which can be a very important consideration when dealing with substrates that are easily fluorinated or oxygenated.

Ongoing studies in our laboratory show that the potential hazards of handling neat HN_3 in the synthesis of N_5^+ can be avoided by either replacing HN_3 with the insensitive $(CH_3)_3SiN_3$ or generating the desired HN_3 from a weighed amount NaN_3 and excess HF in a separate ampoule and transferring all volatiles into the reaction vessel containing an HF solution of $N_2F^+SbF_6$. The reactions with $(CH_3)_3SiN_3$ are carried out in either HF or SO_2 solution and produce N_5^+ in high yield. When HF is used as the solvent, the first reaction step most certainly involves the formation of $(CH_3)_3SiF$ and HN_3 , i. e., HN_3 is generated in situ in the reactor.

Synthesis and Properties of $N_5^+Sb_2F_{11}^-$. To preclude a potential side reaction of $Sb_2F_{11}^-$ with HF and HN₃ to give SbF_6^- and $H_2N_3^+SbF_6^-$, a sample of $N_5^+SbF_6^-$ was reacted with an equimolar amount of SbF_5 in HF solution at room temperature.

$$N_5^+SbF_6^- + SbF_5 \longrightarrow N_5^+Sb_2F_{11}^-$$

The resulting $N_5^+Sb_2F_{11}^-$ salt is a colorless solid that is stable at room temperature and undergoes, according to its DSC data, thermal decomposition at 70 °C, *i. e.*, its thermal stability is comparable to that of $N_5^+SbF_6^-$ but, contrary to $N_5^+SbF_6^-$, it undergoes a reversible endotherm (melting) at about 30 °C. Consequently, the replacement of SbF_6^- by $Sb_2F_{11}^-$ did not result in increased thermal stability and does not appear to offer any significant advantages for studying the reaction chemistry of N_5^+ salts.

Crystal Structure of $N_5^+Sb_2F_{11}^-$. The structure of $N_5^+Sb_2F_{11}^-$ is shown in Fig. 1-4 and the important bond lengths and angles are summarized in Table 3. The observed V-shaped geometry of the N_5^+ cation is in excellent agreement with the theoretical predictions for the free gaseous N_5^+ cation at the B3LYP level of theory with the calculated terminal and central N-N bond distances of 1.11 and 1.30 Å being close to the observed ones of 1.105(19) and 1.299(19) Å, respectively. Furthermore, the terminal N-N distance of 1.105(19) Å in N_5^+ is only slightly longer than that of 1.089(9) Å found for N_2F^+ in $N_2F^+Sb_2F_{11}^-$, and compares well with the N-N bond distances of 1.0976(2) Å in N_2^{-20} and 1.0927 Å found in HN_2^+ , indicating that these bonds approximate triple bonds. The central N-N bond length of 1.299(19) Å in N_5^+ is somewhat longer than those found for typical N-N double bonds (1.17 to 1.25 Å), but is significantly shorter than those found for typical N-N single bonds (1.43 to 1.75 Å). Also, the agreement between

calculated [112.3 and 166.7°] and observed [111.2(11) and 167.2(15)°, respectively] bond angles is very good.

The observed geometry supports the previously given rationale¹ that the exceptional stability of N_5^+ is largely due to resonance stabilization, resulting in relatively high bond orders for all the bonds. The bonding in N_5^+ can be described by the following three resonance structures:

Although the relative contributions from (I) and the equivalent pair, (II) and (III), are unknown, the calculated charge distributions at the NBO(B3LYP/aug-cc-pVDZ) level²⁴

and the relative shielding of the N-nmr signals (the shielding increases from the terminal nitrogen to the β -nitrogen to the central nitrogen in accord with the calculated charge distributions)¹ strongly indicate that the terminal and the β -nitrogen atoms carry the positive charges and the central nitrogen a small negative one.

A least-squares-plane analysis for N_5^+ shows that the cation is essentially planar. The N2 atom exhibits a maximum deviation of 0.11 Å from the average mean plane that shows a root mean square deviation of 0.0058 Å. The N_5^+ mean plane is almost

perpendicular (78.1°) to the plane containing the F5-Sb1-F6-Sb2-F11 atoms. The latter is also almost perfectly planar and shows a root mean square deviation of 0.014 Å.

The geometry of the Sb₂F₁₁ anion also deserves a special comment. This anion is known to possess little rigidity and can exist in either an eclipsed or staggered conformation and exhibit a wide range of Sb-F-Sb bridge angles, depending upon the counter ion present in the crystal lattice.²⁵ The eclipsed conformation is rare but has previously also been observed for BrF₄+Sb₂F₁₁. In the latter compound, the eclipsed structure results from a packing effect in which one equatorial fluorine ligand of each antimony atom of Sb₂F₁₁ bridges to a different BrF₄ cation. Since the two BrF₄ cations and the Sb₂F₁₁ anion are coplanar, the bridging equatorial fluorine ligands around the antimonies become also coplanar resulting in the eclipsed configuration. The eclipsed conformation of the Sb₂F₁₁ anion found for N₅⁺Sb₂F₁₁ (Fig. 2) is also due to fluorine bridging but results from N₅⁺ acting as a spacer between the two equatorial SbF₄ units of Sb₂F₁₁ (Fig. 3). In accord with the resonance structures and the calculated charge distributions of N₅⁺ (see above), the positively charged β-nitrogen atoms interact with the negatively charged fluorine ligands forming bridges that are shorter than the sum of the van der Waals radii of 3.0 Å (Figures 2-4). Thus, N2 and N4 straddle the two eclipsed fluorine atoms, F2 and F8, but since the N₅⁺ plane is not perfectly perpendicular to the F_{ax} -Sb-F-Sb- F_{ax} plane, the N2-F8 = 2.723(15) Å and N4-F2 = 2.887(16) Å distances are somewhat shorter than those of N2-F2 = 3.032(16) Å and N4-F8 = 2.993(16) Å. In addition, N2 and N4 bridge to two fluorine atoms from other Sb₂F₁₁ anions with N4-F4B = 2.813(15) Å and N2-F10A = 2.768(15) Å. In contrast to $BrF_4^+Sb_2F_{11}^-$, which has an

almost linear Sb-F-Sb bridge angle of 175° , 19 that of $155.0(4)^{\circ}$ in $N_5^+Sb_2F_{11}^-$ is much closer to those usually found for $Sb_2F_{11}^{-25}$

Vibrational Spectra of N_5^+ . The infrared and Raman spectra of solid $N_5^+SbF_6^-$ are shown in Figures 5 and 6, respectively. The experimentally observed frequencies of $N_5^+SbF_6^-$, $N_5^+Sb_2F_{11}^-$, and $N_5^+AsF_6^-$, together with their assignments, are listed in Table 4. A comparison of the observed and calculated frequencies and intensities of N_5^+ is given in Table 5. As can be seen, the previously missing remaining 4 fundamental vibrations and numerous combination bands of N_5^+ have been observed and are in excellent agreement with the theoretical predictions for point group C_{2v} . The splittings observed for $v_8(B_2)$ and $v_2(A_1)$ can be accounted for by Fermi resonance. The presence of $Sb_2F_{11}^-$ impurities in the SbF_6^- salt can be readily detected by Raman bands at 692, 598, and 231 cm⁻¹ and infrared bands at 708 and 497 cm⁻¹ that are characteristic for $Sb_2F_{11}^-$ and do not overlap with the SbF_6^- bands.

Conclusion

The synthesis and thorough characterization of $N_5^+SbF_6^-$ and $N_5^+Sb_2F_{11}^-$ demonstrate that the N_5^+ cation can form exceptionally stable salts with fluoroantimonate anions and that these salts are surprisingly insensitive to impact. The N_5^+ cation is a powerful one-electron oxidizer with an electron affinity between 11.0 and 12.08 eV while not giving rise to undesirable fluorination or oxygenation side-reactions. The ready availability of a stable polynitrogen cation in addition to the long known azide anion opens a venue to neutral polynitrogen compounds and may provide the basis for the first synthesis of nitrogen allotropes.

Acknowledgements

The authors thank the Defense Advanced Research Project Agency, the U. S. Air Force Office of Scientific Research, and the National Science Foundation for financial support and Drs. T. Schroer, S. Schneider, N. Maggiarosa, and M. Gerken from the University of Southern California for experimental support and stimulating discussions.

Supporting Information Available

Tables of structure determination summary, atomic coordinates, bond lengths and angles and anisotropic displacement parameters of $N_5Sb_2F_{11}$ in CIF format. This material is available free of charge via the internet at http://pubs.acs.org.

References

- (1) Christe, K. O.; Wilson, W. W.; Sheehy, J. A.; Boatz, J. A. Angew. Chem., Int. Ed. Engl. 1999, 38, 2004.
- (2) See, for example, Chem. Eng, News, Jan. 25, 1999, pg.7; Nov. 29, 1999, pg.38;Aug. 14, 2000, pg. 41.
- (3) Christe, K. O.; Wilson, W. W.; Schack, C. J.; Wilson, R. D. Inorg. Synth.1986, 24, 39.
- (4) Ruff, J. K. Inorg. Chem. 1966, 5, 1971.
- (5) Roesky, H. W.; Glemser, O.; Bormann, D. Chem. Ber. 1966, 99, 1589.
- (6) Pankratov, V. A.; Savenkova, N. I. Zhur. Neorg. Khim. 1968, 13, 2610.
- (7) Christe, K. O.; Wilson, R. D.; Sawodny, W. J. Mol. Structure 1971, 8, 245.
 Christe, K. O.; Wilson, R. D.; Wilson, W. W.; Bau, R.; Sukumar, S.; Dixon, D. A.
 J. Am. Chem. Soc. 1991, 113, 1991.
- (8) Christe, K. O.; Wilson, W. W.; Schack, C. J. J. Fluorine Chem. 1978, 11, 71.
- (9) SMART V 4.045 Software for the CCD Detector System, Bruker AXS, Madison,WI (1999).
- (10) SAINT V 4.035 Software for the CCD Detector System, Bruker AXS, Madison, WI (1999).

[†] Air Force Research Laboratory

[‡] University of Southern California

[§] University of California, Riverside

- (11) SADABS, Program for absorption correction for area detectors, Version 2.01, Bruker AXS, Madison, WI (2000).
- (12) Sheldrick, G. M. SHELXS-97, Program for the Solution of Crystal Structure, University of Göttingen, Germany, 1997.
- (13) Sheldrick, G. M. SHELXL-97, Program for the Refinement of Crystal Structure,
 University of Göttingen, Germany, 1997.
- (14) SHELXTL 5.10 for Windows NT, Program library for Structure Solution and Molecular Graphics, Bruker AXS, Madison, WI (1997).
- (15) Christe, K. O.; Wilson, W. W.; Dixon, D. A.; Khan, S. I.; Bau, R.; Metzenthin, T.;Lu, R. J. Am. Chem. Soc. 1993, 115, 1836.
- (16) Bartlett, N.; Lohmann, D. H.; Proc. Chem. Soc. (London) 1962, 277; J. Chem.
 Soc. 1962, 5253.
- (17) Stein, L.; Norris, J. R.; Downs, A. J.; Minihan, A. R. J. Chem. Soc. Chem. Commun. 1978, 502.
- (18) Stein, L.; Henderson, W. W. J. Am. Chem. Soc. 1980, 102, 2856.
- (19) Vij, A.; Vij, V.; Tham, F.; Christe, K. O. unpublished results.
- (20) Huber, K. P.; Herzberg, G. Constants of Diatomic Molecules; Van Nostrand Reinhold: New York, 1979.
- (21) Owrutsky, J. C.; Gudeman, C. S.; Martner, C. C.; Tack, L. M.; Rosenbaum, N. H.;
 Saykally, R. J. J. Phys. Chem. 1986, 84, 605.
- (22) Botschwina, P. Chem. Phys. Lett. 1984, 107, 535.
- (23) Greenwood, N. N.; Earnshaw, A. Chemistry of the Elements, 2nd ed.;
 Butterworth, Heinemann: Oxford, 1998.

- (24) Fau, S.; Bartlett, R. J. private communication.
- (25) Aubke, F.; Willner, H. Paper C3, presented at the 16th International Symposium on Fluorine Chemistry, Durham, U.K., July 15-20, 2000.

Table 1. Crystal data and structure refinement for $N_5^+\!Sb_2F_{11}^-$

Table 1. Crystal data and structure for	inement for 145 Sb2F1	1
Identification code	$N_5^+Sb_2F_{11}^-$	V.,
Empirical formula	F11 N5 Sb2	
Formula weight	522.55	
Temperature	213(2) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	C2/c	•
Unit cell dimensions	a = 10.913(8) Å	L → 1 = 90°.
	b = 12.654(8) Å	/3 Æ 104.715(18)°.
	c = 16.675(11) Å	و 90° = 90°.
Volume	2227(3) Å ³	
Z	8	
Density (calculated)	3.117 Mg/m ³	
Absorption coefficient	4.995 mm ⁻¹	
F(000)	1888	
Crystal size	0.26 x 0.10 x 0.05 mi	m^3
Theta range for data collection	2.51 to 25.35°.	
Index ranges	-12<=h<=13, -15<=k	<=15, -20<=l<=17
Reflections collected	9125	
Independent reflections	2022 [R(int) = 0.0629]	9]
Absorption correction	Sadabs	
Max. and min. transmission	0.7883 and 0.3567	
Refinement method	Full-matrix least-squa	ares on F ²
Data / restraints / parameters	2022 / 0 / 164	
Goodness-of-fit on F ²	1.122	
Final R indices [I>2sigma(I)]	R1 = 0.0678, $wR2 = 0.0678$	0.1913
R indices (all data)	R1 = 0.0785, $wR2 = 0$	0.2019
Extinction coefficient	0.00026(18)	
Largest diff. peak and hole	4.329 and -2.102 e.Å	-3

Table 2. Atomic coordinates (x 10^4) and equivalent isotropic displacement parameters (Å 2 x 10^3) for $N_5^+Sb_2F_{11}^-$. U(eq) is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	X	у	Z	U(eq)	
Sb(1)	2327(1)	5268(1)	5963(1)	32(1)	
Sb(2)	1229(1)	2343(1)	6199(1)	28(1)	
F(1)	930(11)	5546(9)	6373(7)	65(3)	
F(2)	3187(10)	4839(8)	7017(5)	59(3)	
F(3)	3652(9)	4765(7)	5561(6)	48(2)	
F(4)	1368(9)	5453(7)	4889(5)	51(2)	
F(5)	2906(11)	6638(7)	6101(6)	64(3)	
F(6)	1691(8)	3755(6)	5813(5)	43(2)	
F(7)	-17(9)	3062(8)	6548(6)	53(2)	
F(8)	2340(9)	2672(7)	7217(5)	50(2)	
F(9)	2592(10)	1837(9)	5844(6)	63(3)	
F(10)	190(10)	2218(6)	5142(5)	49(2)	
F(11)	854(10)	1003(6)	6539(6)	59(3)	
N(1)	4750(14)	3132(16)	8389(9)	62(4)	
N(2)	3965(12)	3518(11)	8581(7)	40(3)	
N(3)	3138(14)	3881(10)	8954(8)	51(4)	
N(4)	2229(13)	4376(10)	8441(7)	40(3)	
N(5)	1369(14)	4828(13)	8136(8)	52(4)	

Table 3. Bond lengths and angles for $N_5^+Sb_2F_{11}^-$

Bond lengths (Å)

Sb(1)-F(5)	1.839(8)	Sb(2)-F(9)	1.849(10)
Sb(1)-F(4)	1.845(8)	Sb(2)-F(8)	1.866(8)
Sb(1)-F(2)	1.851(9)	Sb(2)-F(11)	1.866(8)
Sb(1)-F(3)	1.854(9)	Sb(2)-F(6)	2.007(7)
Sb(1)-F(1)	1.856(10)	N(1)-N(2)	1.102(19)
Sb(1)-F(6)	2.031(7)	N(2)-N(3)	1.303(19)
Sb(2)-F(10)	1.844(7)	N(3)-N(4)	1.295(19)
Sb(2)-F(7)	1.849(9)	N(4)-N(5)	1.107(19)
Bond Angles (°)			
F(5)-Sb(1)-F(4)	95.4(4)	F(9)-Sb(2)-F(8)	88.8(5)
F(5)-Sb(1)-F(2)	94.8(4)	F(10)-Sb(2)-F(11)	94.8(4)
F(4)-Sb(1)-F(2)	169.9(4)	F(7)-Sb(2)-F(11)	96.7(5)
F(5)-Sb(1)-F(3)	95.4(5)	F(9)-Sb(2)-F(11)	92.2(5)
F(4)-Sb(1)-F(3)	89.7(4)	F(8)-Sb(2)-F(11)	93.9(4)
F(2)-Sb(1)-F(3)	89.3(5)	F(10)-Sb(2)-F(6)	85.0(3)
F(5)-Sb(1)-F(1)	93.6(5)	F(7)-Sb(2)-F(6)	85.9(4)
F(4)-Sb(1)-F(1)	91.0(5)	F(9)-Sb(2)-F(6)	85.3(5)
F(2)-Sb(1)-F(1)	88.4(5)	F(8)-Sb(2)-F(6)	86.3(4)
F(3)-Sb(1)-F(1)	170.8(4)	F(11)-Sb(2)-F(6)	177.5(4)
F(5)-Sb(1)-F(6)	179.9(5)	Sb(2)-F(6)-Sb(1)	155.0(4)
F(4)-Sb(1)-F(6)	84.7(4)	N(1)-N(2)-N(3)	168.1(15)
F(2)-Sb(1)-F(6)	85.1(4)	N(4)-N(3)-N(2)	111.2(11)
F(3)-Sb(1)-F(6)	84.5(4)	N(5)-N(4)-N(3)	166.3(14)
F(1)-Sb(1)-F(6)	86.4(4)		
F(10)-Sb(2)-F(7)	91.2(5)		
F(10)-Sb(2)-F(9)	90.8(5)		
F(7)-Sb(2)-F(9)	170.7(5)		
F(10)-Sb(2)-F(8)	171.3(4)		
F(7)-Sb(2)-F(8)	87.9(4)		

Table 4. Observed Infrared and Raman Spectra of $N_5^+SbF_6$, $N_5^+Sb_2F_{11}^-$ and $N_5^+AsF_6^-$ and

Their Assignments

-----assgnts (point group)-----assgnts (point group)--------- $N_5^+Sb_2F_{11}^-$ ---- $N_5^+AsF_6^-$ --- $MF_6(O_h)$ Sb_2F_{11} $N_5^+(C_{2v})$ IR RA IR RA IR RA $(v_1 + v_3 + v_9)(B_2) =$ 3357 3358 vw $(v_1 + v_8)(B_2) = 3323$ 3334 vw 3069 w $(v_2 + v_7)(B_2) = 3077$ 3079 w $(v_1 + v_9)(B_2) = 2682$ 2671 vw 2681 vw 2261 (9.0) 2270 2271 (4.4) $(v_1)(A_1)$ 2270 m 2268 2260 m (9.4)2202 (1.9) 2210 s $(v_7)(B_2)$ 2211 (0.8) 2205 2203 s 2205 s (2.0) $(v_3 + 3v_9)(B_2) = 1914$ 1921 1919 vw vw $(v_8 + 2v_9)(B_2) = 1883$ 1891 1883 vw vw Comb. Comb. 1366 w 1240 bands bands 1288 vw[٧W $(v_3 + v_9)(B_2) = 1086^a$ 1088 s 1092 1089 s ms $v_8(B_2)$ 1064 s 1064 s $(v_5 + v_6)(B_2) = 903$ 902 892 vvw vvw $v_2(A_1)$ 866 (0.6) 872 w 871 (0.7) 871 w 872 (0.6) 867 w $(2v_9)(A_1) = 828^b$ 824(0+)835 vw 837 (0+) 824 vw 692 (5.5) 725 vSbF 654 (10) 650 vs,br 598 (1.4) 669/672 $v_3(A_1)$ 680 sh 664 (~1) 672 (1) (1.8) $v_3(F_{1u})$ 655 vs 704 vs $\nu_l(A_{lg})$ 686 (10) 652 (10) 596 mw vSbF 537 mw $v_2(E_g)$ 575 w 579 (1.6) 582 w 571 (0.8) vSb-F-Sb 497 s 478 (0+) 470 (0+) $v_5(A_2)$ 447 w 449 w 420 sh $v_6(B_1)$ 417 ms 425 ms 417 (0+) $v_0(B_2)$ 412 mw 416 (0+) 409 sh 295 (2.1) 283 sh δSb-F 272 sh 231 (2.0) $v_4(F_{1u})$ 394 vs 284 vs $v_5(F_{2g})$ 372 (3.4) 282 (2.8) 204 (5.0) 200 (3.6) 209 (4.4) $\nu_4(A_1)$ 135 sh 125 (5.5) lattice vibrations 107 (5) 97 (5.0)

a) In Fermi resonance with $v_8(B_2)$. b) In Fermi resonance with $v_2(A_1)$

Table 5. Comparison of Observed and Unscaled Calculated CCSD(T)/6311+G(2d) Vibrational Frequencies (cm $^{\text{-}1}$) and Intensities (km mol $^{\text{-}1}$ and $\text{Å}^{\text{-}4}$ amu $^{\text{-}1}$) for $N_5^{\text{+}}$

	approx mode description in point group C _{2v}	calcd fro	eq (abs IR	R, Ra int)	obsd freq (re	el IR, RA	(int)
A_1	v_1 in-phase terminal stretches	2229	(13)	[215]	2260-2271	m	[10.0]
	v_2 sym central stretch	818	(0.5)	[5]	866-872	w	[0.6]
	v ₃ central bending	644	(2)	[1]	664-672	obscd	[1] ^a
	v ₄ in-phase terminal bends	181	(0.3)	[6]	200-209	_b	[4]
A ₂	v_5 out-of-phase, out-of-plane bend	475	(0)	[1]	470-478	-	[0+]
B_1	ν ₆ in-phase, out-of-plane bend	405	(6)	[0]	417-425	ms	[0]
B_2	ν ₇ out-of-phase term stretches	2175	(105)	[42]	2203-2211	S	[1.9]
	v ₈ asym central stretch	1032	(138)	[2]	1055-1064	s .	[n. obsd]
	ν ₉ out-of-phase term bends	399	(1)	[0.5]	412-417	mw	[+0]

^aObscured in infrared and interference in Ra by anion bands. ^bOutside of frequency range of our spectrometer.

Figure Captions

- Figure 1. An ORTEP diagram of N_5 *Sb₂F₁₁ showing the thermal ellipsoids at the 30% probability level.
- Figure 2. An ORTEP diagram showing the side view of the eclipsed Sb_2F_{11} anion. The thermal ellipsoids are drawn at the 30% probability level.
- Figure 3. Space filling representation of $N_5^+Sb_2F_{11}^-$ showing the close packing of the N_5^+ cation within the $Sb_2F_{11}^-$ cavity.
- Figure 4. A figure showing close range N···F contacts within the crystal lattice of N_5 ⁺Sb₂F₁₁⁻.
- Figure 5. Infrared spectrum of solid $N_5^+SbF_6^-$ recorded as an AgBr pellet at room temperature. The band at 498 cm⁻¹, marked by an asterisk, is due to a small amount of $Sb_2F_{11}^-$.
- Figure 6. Raman spectrum of solid $N_5^+SbF_6^-$ recorded at room temperature.





Figure 1






Figure 2




Figure 3





Figure 4



SUPPLEMENTARY MATERIAL

Polynitrogen Chemistry. Synthesis, Characterization, and Crystal Structure of Surprisingly Stable Fluoroantimonate Salts of ${\rm N_5}^+$

Ashwani Vij, *William W. Wilson, *Vandana Vij, *Fook S. Tham, *Jeffrey A. Sheehy, *and Karl O. Christe, ***

Propulsion Sciences and Advanced Concepts Division, Air Force Research Laboratory (AFRL/PRS), Edwards AFB, California 93524, Loker Hydrocarbon Research Institute and Department of Chemistry, University of Southern California, Los Angeles, CA 90089, and Department of Chemistry, University of California, Riverside, CA 92521

Abstract

The new N_5^+ salt, N_5^+ SbF₆, was prepared from N_2F^+ SbF₆ and HN₃ in anhydrous HF solution. The white solid is surprisingly stable, decomposing only at 70 °C, and is relatively insensitive to impact. Its vibrational spectrum exhibits all nine fundamentals with frequencies that are in excellent agreement with the theoretical calculations for a five-atomic V-shaped ion of C_{2v} symmetry. The N_5^+ Sb₂F₁₁ salt was also prepared and its crystal structure was determined. The geometry previously predicted for free gaseous N_5^+ from theoretical calculations was confirmed within experimental error. The Sb₂F₁₁ anions exhibit an unusual geometry with eclipsed SbF₄ groups due to inter-ionic bridging with the N_5^+ cations. The N_5^+ cation is a powerful one-electron oxidizer. Its electron affinity falls between 11.0 and 12.08 eV because it readily oxidizes NO to NO⁺ and NO₂ to NO₂⁺ but fails to oxidize Xe or O₂.

Courted data and structure refinement for N5SB2F11.

Largest diff. peak and hole

Table 1. Crystal data and structure refinement	ent for N5SB2F11.	
Identification code	N ₅ Sb ₂ F ₁₁	
Empirical formula	F11 N5 Sb2	
Formula weight	522.55	
Temperature	213(2) K	
Wavelength	0.71073 Å	•
Crystal system	Monoclinic	
Space group	C2/c	α= 90°.
Unit cell dimensions	a = 10.913(8) Å	$\beta = 104.72(2)^{\circ}$.
	b = 12.654(8) Å	$y = 90^{\circ}$.
	c = 16.675(11) Å	$\gamma = 90^{\circ}$.
Volume	2227(3) Å ³	•
Z	8	
Density (calculated)	3.117 Mg/m^3	
Absorption coefficient	4.995 mm ⁻¹	
F(000)	1888	
Crystal size	$0.26 \times 0.10 \times 0.05 \text{ mm}^3$	·
Theta range for data collection	2.51 to 25.35°.	
Index ranges	-12<=h<=13, -15<=k<=	15, -20<=1<=17
Reflections collected	9125	
Independent reflections	2022 [R(int) = 0.0629]	
Completeness to theta = 25.35°	99.6 %	
•	SADABS	
Absorption correction Max. and min. transmission	0.7883 and 0.3567	
	Full-matrix least-square	es on F ²
Refinement method	2022 / 0 / 164	
Data / restraints / parameters	1.122	
Goodness-of-fit on F ²	R1 = 0.0678, $wR2 = 0$.	1913
Final R indices [I>2sigma(I)]	R1 = 0.0785, $wR2 = 0$	
R indices (all data)	0.00026(18)	
Extinction coefficient	4.329 and -2.102 e.Å-3	
Largest diff. peak and hole		

Table 2. Atomic coordinates (x 10⁴) and equivalent isotropic displacement parameters (Å²x 10³) for N5SB2F11. U(eq) is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	· x	у	z	U(eq)
Sb(1)	2327(1)	5268(1)	5963(1)	32(1)
Sb(2)	1229(1)	2343(1)	6199(1)	28(1)
F(1)	930(11)	5546(9)	6373(7)	65(3)
F(2)	3187(10)	4839(8)	7017(5)	59(3)
F(3)	3652(9)	4765(7)	5561(6)	48(2)
F(4)	1368(9)	5453(7)	4889(5)	51(2)
F(5)	2906(11)	6638(7)	6101(6)	64(3)
F(6)	1691(8)	3755(6)	5813(5)	43(2)
F(7)	-17(9)	3062(8)	6548(6)	53(2)
F(8)	2340(9)	2672(7)	7217(5)	50(2)
F(9)	2592(10)	1837(9)	5844(6)	63(3)
F(10)	190(10)	2218(6)	5142(5)	49(2)
F(11)	854(10)	1003(6)	6539(6)	59(3)
N(1)	4750(14)	3132(16)	8389(9)	62(4)
N(2)	3965(12)	3518(11)	8581(7)	40(3)
N(3)	3138(14)	3881(10)	8954(8)	51(4)
N(4)	2229(13)	4376(10)	8441(7)	40(3)
N(5)	1369(14)	4828(13)	8136(8)	52(4)

Table 3. Bond lengths [Å] and angles [°] for N5SB2F11.

Sb(1)-F(5)	1.839(8)
Sb(1)-F(4)	1.845(8)
Sb(1)-F(2)	1.851(9)
Sb(1)-F(3)	1.854(9)
Sb(1)-F(1)	1.856(10)
Sb(1)-F(6)	2.031(7)
Sb(2)-F(10)	1.844(7)
Sb(2)-F(7)	1.849(9)
Sb(2)-F(9)	1.849(10)
Sb(2)-F(8)	1.866(8)
Sb(2)-F(11)	1.866(8)
Sb(2)-F(6)	2.007(7)
N(1)-N(2)	1.102(19)
N(2)-N(3)	1.303(19)
N(3)-N(4)	1.295(19)
N(4)-N(5)	1.107(19)
F(5)-Sb(1)-F(4)	95.4(4)
F(5)-Sb(1)-F(2)	94.8(4)
F(4)-Sb(1)-F(2)	169.9(4)
F(5)-Sb(1)-F(3)	95.4(5)
F(4)-Sb(1)-F(3)	89.7(4)
F(2)-Sb(1)-F(3)	89.3(5)
F(5)-Sb(1)-F(1)	93.6(5)
F(4)-Sb(1)-F(1)	91.0(5)
F(2)-Sb(1)-F(1)	88.4(5)
F(3)-Sb(1)-F(1)	170.8(4)
F(5)-Sb(1)-F(6)	179.9(5)
F(4)-Sb(1)-F(6)	84.7(4)
F(2)-Sb(1)-F(6)	85.1(4)
F(3)-Sb(1)-F(6)	84.5(4)
F(1)-Sb(1)-F(6)	86.4(4)
F(10)-Sb(2)-F(7)	91.2(5)
F(10)-Sb(2)-F(9)	90.8(5)
* .	

F(7)-Sb(2)-F(9)	170.7(5)
F(10)-Sb(2)-F(8)	171.3(4)
F(7)-Sb(2)-F(8)	87.9(4)
F(9)-Sb(2)-F(8)	88.8(5)
F(10)-Sb(2)-F(11)	94.8(4)
F(7)-Sb(2)-F(11)	96.7(5)
F(9)-Sb(2)-F(11)	92.2(5)
F(8)-Sb(2)-F(11)	93.9(4)
F(10)-Sb(2)-F(6)	85.0(3)
F(7)-Sb(2)-F(6)	85.9(4)
F(9)-Sb(2)-F(6)	85.3(5)
F(8)-Sb(2)-F(6)	86.3(4)
F(11)-Sb(2)-F(6)	177.5(4)
Sb(2)-F(6)-Sb(1)	155.0(4)
N(1)-N(2)-N(3)	168.1(15)
N(4)-N(3)-N(2)	111.2(11)
N(5)-N(4)-N(3)	166.3(14)
• • •	

Symmetry transformations used to generate equivalent atoms.

Table 4. Anisotropic displacement parameters (Å 2 x 10 3) for N5SB2F11. The anisotropic displacement factor exponent takes the form: $-2\pi^2$ [h^2 a^{*2} U 11 + ... + 2 h k a^* b* U 12]

	Ω_{11}	U_{55}	Ω_{33}	U^{23}	Ω_{13}	U_{15}
Sb(1)	46(1)	20(1)	33(1)	-4(1)	16(1)	-10(1)
Sb(2)	37(1)	15(1)	29(1)	0(1)	2(1)	-3(1)
F(1)	80(7)	58(6)	74(7)	-15(5)	48(6)	1(5)
F(2)	73(6)	69(7)	27(4)	2(4)	0(4)	-37(5)
F(3)	48(5)	50(6)	49(5)	-4(4)	19(4)	-5(4)
F(4)	64(6)	43(5)	41(5)	14(4)	5(4)	-8(4)
F(5)	101(8)	16(4)	73(7)	-10(4)	21(6)	-14(5)
F(6)	64(5)	29(4)	35(4)	1(3)	10(4)	-21(4)
F(7)	63(6)	48(6)	54(5)	0(4)	26(4)	10(5)
F(8)	63(6)	45(5)	30(4)	-1(4)	-12(4)	-6(4)
F(9)	52(6)	69(7)	70(6)	-14(5)	19(5)	15(5)
F(10)	78(6)	29(4)	25(4)	-2(3)	-16(4)	-7(4)
F(11)	80(7)	24(5)	61(6)	13(4)	-2(5)	-7(4)
N(1)	43(8)	103(14)	40(7)	-3(8)	8(6)	8(8)
N(2)	45(7)	46(7)	23(5)	0(5)	-1(5)	-6(6)
N(3)	77(10)	40(8)	39(7)	0(6)	19(7)	22(7)
N(4)	54(8)	37(7)	33(6)	-2(5)	18(6)	-12(6)
N(5)	51(8)	68(10)	41(7)	-1(7)	18(6)	-2(7)

Table 5. Torsion angles [°] for N5SB2F11.

F(10)-Sb(2)-F(6)-Sb(1)	156.0(13)
F(7)-Sb(2)-F(6)-Sb(1)	64.4(13)
F(9)-Sb(2)-F(6)-Sb(1)	-112.8(13)
F(8)-Sb(2)-F(6)-Sb(1)	-23.7(12)
F(11)-Sb(2)-F(6)-Sb(1)	-118(7)
F(5)-Sb(1)-F(6)-Sb(2)	47(100)
F(4)-Sb(1)-F(6)-Sb(2)	-154.6(13)
F(2)-Sb(1)-F(6)-Sb(2)	25.4(12)
F(3)-Sb(1)-F(6)-Sb(2)	115.2(13)
F(1)-Sb(1)-F(6)-Sb(2)	-63.3(13)
N(1)-N(2)-N(3)-N(4)	-173(8)
N(2)-N(3)-N(4)-N(5)	-179(100)

Symmetry transformations used to generate equivalent atoms.

Table 6 - Contact Distances(Angstrom) for N5SB2F11

- Sb1 .N3_a 3.830(14) F8 .F2 2.940(14)
- Sb1 .F11_b 3.837(11) F8 .N1 2.909(18)
- Sb2 .F5_c 3.697(12) F8 .N2 2.723(15)
- F1 .N5 2.997(18) F8 .N3 3.195(15)
- F1 .N5_d 2.97(2) F8 .N4 2.993(15)
- F1 .F4_e 3.104(15) F9 .N5_j 3.108(19)
- F2 .F8 2.940(14) F9 .F3_h 3.135(14)
- F2 .N2 3.032(15) F10 .N1_k 2.874(17)
- F2 .N4 2.887(16) F10 .N2_k 2.768(15)
- F2 .N5 3.049(18) F10 .N3_k 2.936(17)
- F2 .F11_f 2.790(13) F11 .Sb1_c 3.837(11)
- F3 .N1_g 2.97(2) F11 .F3_c 2.983(14)
- F3 .N2_g 3.068(17) F11 .F2_1 2.790(13)
- F3 .N3_a 3.109(16) F11 .N2_j 3.160(16)
- F3 .F11_b 2.983(14) F11 .N3_j 3.091(16)
- F3 .F9_h 3.135(14) F11 .N4_j 2.929(17)
- F4 .F1_e 3.104(15) N1 .F8 2.909(18)
- F4 .F7_e 3.107(13) N1 .N4 3.19(2)
- F4 .N3_a 2.899(18) N1 .F3_m 2.97(2)
- F4 .N4_a 2.813(15) N1 .N1_g 3.15(2)
- F4 .N5_a 2.945(16) N1 .F10_n 2.874(17).
- F5 .Sb2_b 3.697(12) N2 .F2 3.032(15)
- F5 .F7_b 2.843(15) N2 .F8 2.723(15)
- F5 .N3_i 3.051(16) N2 .N5 3.20(2)
- F7 .N4_d 2.935(17) N2 .F3_m 3.068(17)
- F7 .N5_d 2.801(19) N2 .F11_f 3.160(16)
- F7 .F4_e 3.107(13) N2 .F10_n 2.768(14)

```
3.195(15)
               2.843(15) N3
                                .F8
       .F5_c
∠ F7
                                         2.813(15)
                3.830(14) N4
                                 .F4_o
        .Sb1_o
  N3
                                .F11_f
                                       2.929(17)
                           N4
                3.109(16)
  N3
        .F3_o
                                .F1
                                       2.997(18)
                           N5
                2.899(18)
   N3
        .F4_o
                                       3.049(18)
                                .F2
                          N5
                3.051(16)
        .F5_1
   N3
                                         3.20(2)
                          N5
                                 .N2
                3.091(16)
        .F11_f
   N3
                                          2.97(2)
        .F10_n 2.936(17) N5
                                 .F1_p
   N3
                                       2.801(19)
               2.887(16) N5
                                .F7_p
        .F2
   N4
                                .N5_d
                                         3.19(2)
               2.993(15)
                          N5
   N4
        .F8
                                       2.945(16)
                 3.19(2) N5
                               .F4_o
   N4
        .N1
                                        3.108(19)
                                .F9 f
                2.935(17) N5
         .F7_p
```

Translation of Symmetry Code to Equiv.Pos

```
a = [4564.00] = x,1-y,-1/2+z
b = [5555.00] = 1/2 + x, 1/2 + y, z
c = [5445.00] = -1/2+x, -1/2+y, z
d = [2556.00] = -x,y,3/2-z
e = [3566.00] = -x,1-y,1-z
f = [6556.00] = 1/2-x, 1/2+y, 3/2-z
g = [2656.00] = 1-x,y,3/2-z
h = [7556.00] = 1/2-x, 1/2-y, 1-z
i = [6556.00] = 1/2-x,1/2+y,3/2-z
j = [6546.00] = 1/2-x,-1/2+y,3/2-z
k = [8454.00] = -1/2+x, 1/2-y, -1/2+z
1 = [6546.00] = 1/2-x, -1/2+y, 3/2-z
m = [2656.00] = 1-x,y,3/2-z
n = [8555.00] = 1/2+x,1/2-y,1/2+z
o = [4565.00] = x,1-y,1/2+z
p = [2556.00] = -x, y, 3/2-z
```

Table

																															-					_						
s	S	6	3	_		7 ,		σ.	4	۲.	4	ж - с	٠, ٠	T 0	۰ د	J <	r (V 5	Jr (, م	7 1	ഹം	5 0	ۍ د	1 0	o L	. ,		. ~	. 2	v	, m	٠,	, ,		٠	, r	۰ -	, -	7 _	- a	o
c 10	3	3 1																																								0 70
10Fc	115	153	630	40	100	86.	1	25	98	54	49	4. 2. c	7 0	0 0	, ,	200	7 0	132	7 C T	9 6	, כ	4	- כ	2 6	, <	÷ ~	2	76	11	88	25	284	17	320	00	248	5	226	10.7	47	200	80
10F0	110	542	669	466	.013	906	123	90	6/8	617	548	212	ָ ק	411	9 6	707	000	470.	0 0	907	0 1	273	1 0	247	- 000	988	0	843	0	005	289	1756	185	052	883	182	77	196	2	461	230	20
7 7	2 1	7	7	7	7 7	2 (7 (2 (7	2	~ (7 (٦ ,	4 0	1 0	4 C	1 0	4 6	4 (, c	1 (7 c	1 0	7 C	10	۷ ۷	. ~	m	m	3	٣	3	m	(1)	٠,	, ~		י נ) (L	י ר		າຕ
×	11	11	11	77	7 ;	- ;	Ţ.	12	7 7	7.7	7.7	77	1 5	1 0	1 .	3 .	, ,	3 5	, ,	17	7 -	7 -	, ,	7 -	٦,	12	15	-	Н	7	Н	-	~	٦	-		۱	4	٠,	٠,-	۱۸	1 7
4	-5		-1	~ (יי	Ωį	- (ap (۰ و	T (7 .	c د	1 <	r (c	ים ו	ן מ	1		4 0	ט ע	,	1 1	10	o ر	1 4	- 1	-	-13	-11	6-	1	-5		-1	-	~ ۱	ı ıc	, ,	- σ	١, ٢	-12	-10
10s	56	25	58	56	77	16	010	7.7	٦ ,	, d	4 g	77	F 7	25] [15	9 4	22	1 0	30	י ע נ		1,	25	5.0	14	32	24	15	47	45	43	54	09	18	20	42	10	17	20	36	34
ည	482	026	116	562	1/1	9 0	0 0	248	130	797	910	707	י כ ני כ	118	200	256	442	440	700	280	. 000	900	712	2275	270	513	984	062	441	242	252	410	654	37	222	498	970	399	735	768	164	320
o 10																																										
10F0	243	142	461		8 6	70,	4 4	ດິ	ć	ח ל	0 1	7 2	,	23	8	122	77	4	1 4		2	ó	Ö	3175	346	9	217	212	14]	15	7	4(6	v	132	9	426	4	183	16	246	2,
7	3	3	2 0		9 0	, ,	7 6	2 6	, ,	, c	, ,	, 0	1 6	. 6	2	. 4	2	1 2				1 0		10	5 2	2	5 2	5 2	5	5 2	5.2	5 2	5	5 2	5 2	5 2	5	2	5	.0	5	5 2
ч	ώ 	ņ.		٠.	ი <u>.</u>	ט ר	- 0	ν -		, <	٥ د	ې د	, 4	٠,	0	2	4	. 9	α	0		. 6		- rċ	٠,	-		د	r,	7	σ.	Ξ.	_ O	- 80	و	4	.5	0	7	4	9	8
		•	1					Ť	7 (7											ī	٠,	,	- 5	,	,							ī	'	•	,	1					
10s	~	~	31	7,7		200	2 6	7 7	, 5	3 6	7 6	30	34	24	40	18	29	24	40	31	32	42	19	19	21	22	29	20	26	122	42	22	22	70	74	20	21	57	27	30	92	36
10Fc	143																							1682																		
OFo :	0																							1670																		
1 10	٦,		, c	, ,	1.6	4	-	7 -	, ,	,	, <u>,</u>	; ~	-	1.2	1 19	1 11		-	7	, , ,	7	1 10	1 11	1 16	1 16	7	7	1. 1.	1 21	7		1 16	1 1	7	.,	1 10	1 11	1 6	1 11	1 12	7	1 16
×	80 4	э (ט פ	ησ	٠ ٥	٠ ٥	٠ ٥	0	, o	٠ ٥	, 5	10	10	10	10	10	10	10	10	11	11	11	11	11	11	1.1	11	12	12	12	12	12	12	12	13	13	13	13	13	13	1.4	14
ح	10	ן טינ	- 4	י ר	, -		. ~	י ע	, ,	- σ	ν α 1	9	4-4	-2	0	7	4	9	8	-7	.5	6-	7	7	m	2	7	9-	-4	-2	0	2	4	9	-5	ű	7	7	m	2	-4	-2
70	ω.	 ,	~ r	- ~		. ~	c		. 57				~	_	0	Ç.	_	7	10			~:	Ψ.	~	_	-	-			m /	m	<u>م</u>	_	~		/	٠.	•	~	٥.		01
: 10s	2,																							13																		
10FC	2084	א לי	1395	146	573	628	646	650	1334	305	5(412	128(101	383	29(1003	215	2086	633	1927	1863	5654	1325	179	1031	1834	3	632	246	ਲ (1705	243	460	290	1460	653	477	843	325	100	762
OFO	750	ם ני	787	130	609	586	675	832	619	403	0	443	147	0	368	197	013	211	670	629	991	221	532	1238	715	957	810	81	670	123	738	687	519	604	694	009	611	486	843	291	0	791
1 1	1 1	٦ -	- -	-, ،		· ~	_	· ~	-	-		7	1 1	_	~	7	7	-	1 2	Н		1 2	1	1 1	~		~ ~	٠, ٠,	<u>ب</u>	,,	٠,	.	7 1	-		1	- -			д	 ,	-
ᅩ	 -	٦ +		1 ~	2	2	7	2	2	7	7	7	7	5	7	7	က	3	m	n	c	3	٣	c	m	ლ (· C	m ·	4	4.	7' '	φ,	7 .	4	4	4	4	4	4	٠ .	σ,	2
r	ស្រ	۰ ۵	y [-12	-10	-8	9-	-4	-2	0	7	4	9	8	10	12	-11	6-	-7	-5	۳-	7	, —(co	S	7	,	11	-12	-10	יס ו	9 •	7 (-2	0	2	4	9	œ	07	7.	-13
10s	88	0 0	106	72	27	10	23	18	17	33	27	44	19	25	16	27	49	29	22	84	16	24	31	419	56	19	32	23	70	n G	0 0	81 .	- 0	28	54	-1	23	26	28	26	2 fr	87
10Fc	4655	3166	252	514	5	774	1246	1141	963	749	310	4048	2369	1189	829	664	756	1900	1339	81	841	1226	788	6174	7.14.7	344	1163	2549	391	263	000	2000	1000	1344	769	108	310	3129	306	529	423	7 2
10Fo	4431	786	219	500	173																																		35	519	000	٦ 8
	0 0																							_															5	> 0	> 0	· >
×	00	> C	0	0	~	٦	~		~	٦	7	2	7	7	7	7	7	m i	က	ო :	m	m ·	m	4.	4	4.	3 7	4 <	. .	ኋ ሊ	י כ	Ωú	ם כ	ກເ	n 1	ა,	ره	y Q	۰ ,	י פ	י ס	-
ڃ	7 4	α	10	12	-	ო	5	7	6	11	0	7	4	9 (œ	10	12	H .	m	ι Ω	1	σ,	7	0 (ν.	4.0	0 0	ם כ	1,0	7 -	٦ ,	י ר	י כ	\	٦,	7	-	4	9 0	α ς	7	4

10s	26 21 19	30	15	18	65	48	33	31	89	21	44 25	19	21	81	81	77	39	16	83	14	21	-	7	67	79	20	, ,	48	2 %	13	17	20
10Fc	321 1457 1299	915 1571 1554	782	822	297	984	1671	1440	50	1746	280	720	196	242	121	609	1768	440	57	1303	6891	96	25	899	233	101	378	854	119	1392	721	2041
.0Fo	331 580 215	694 601 7 524 1	710	174	263	943	109 386 1	552	89	602 1	513	731	943	300	216	648	990 1	483	89	730	537]	0	0	660	707	. 661	. 686	190	. 201	1428 1	693	924
i a	888	~ ~	•		-	-	~	4	,	٦.	-	1					_			-	-											
×	2222									mr		m	٣	4	7,	1 d	. 4.	4	φ,	. 4	٠ 4									2		
ء	8 9 4 (7-0	140	8 01	-11	6-	ر د ا	1-1	, (m u	٠ ٦	6	11	-12	-10	0 Y	-4	-2	0 (N 4	. 0	8	10	-11	ן ן שר	٠ ر٠	י נ	1		·m	2	7
10s	68 1 36	6 / 25 82	34 35	23	20	35	157 21	33	21	56 25	26	24	32	26	و ر	25	39	25	1.9	20	23		41	52	0 7 0	r (C	28	80	29	65	42.	11
10Fc	250 170 985	314	2934 2088	253	720	667	210	274	2007	8/	358	971	389	671	330	992	1607	1527	776	1009	718	80	393	54/	1119	320	1054	149	708	349	559	646
10Fo		371																														
٦	2222																															
*	911		7	7	1	· & c	ဆထ	8	œ c	_∞ α	, ω	8	8	ω (ת ס	0	6	σ.	o 0	ם ע	6	6	10	0 0	200	10	10	10	10	10	11	11
ч	11.	3 22	1-1	და	7	-10	ρ 	-4	-2	2 0	4	9	80	10	ן ן טר	-5	-3	7	r	o ro	7	σ,	φ ₍	0 7	- 2	0	2	4	9	80	6-1	-7
10s	21 24 34				_				42							10		12		1	c									42	31	22
10Fc	1502 656 451	414 615 256	1394 679	571 1732	5801	968	1817	142	540	1416	2248	3709	3359	2743	67.03	281	241	210	759	605	2727	79	9791	832	1267	3795	1395	253	009	175	1.263	2172
FO	1540 678 429																														402	LO.
1 10	20 4 9	2		2 52 2 14																											_	2
٠	4446										.,	· ·	., .	-, -							2											m
r	2 1 2 1 4 1 1	1 1 2	0 8	4 0	2 4	90	0	2 .	д σ	<u>ب</u> ،	ις	ლ,	٠,	~												9	8	0	2	 (თ :	-
	'	1 12 -12							-11		٠,																			-1		
10s	164 17 21 48	24	20	16	36	21	188	82	3 5	16	1.8	39	20 6	25	18	18	19	21	19	38	21	50	63	52	25	59	17	23	18	19	20	2.3
10Fc		1704									96		9 6		122							150		33.6	193	350	66		21	146	ဂ :	2
10Fo	164 1181 391.	337	2255 3034	855 539	582 1105	1245	924	633	1962	1314	928	269	360	232	240	157	345	1054	089	217	410	207	322	282	.987	8688	.251	24	269	491	556	711
-				-			-				-		٦,		-				, –	_	Α,	— — .	- ,	·	7	-		٦,	~ ·		٠,	⊣
×	വവവ	5	5	വവ	9	9 9	و ر	6	و ه	9	9	9	٦ ٥	٠,	7	7	٦ ،	٠ ر	. ~	7	۲ ،	- 1	- &	8	8	æ	ထ	σ,	ω ,	သာင	∞ α	a
ч	9111		നഹ	6	11	8 9	-4	7.5	o 0	4	9	ж <u>г</u>	10	6-1	-7	-5	۲. د	 -	ı m	ທີ	ر ر	٦ ر	-10	8-	9-	-4	-2	0 (7	4 A	ه ه	o
10s	28 29 19 55	52 80	17	42	1 44	18 26	38	30	30	20	42	35	4 t	20	23	56	25	7	20	28	42	, o	49	24	41	48	37	21	41	77	27	71
10Fc	588 783 721 461	463 2123	792	271 899	110 1809	1044 950	570	1342	482	1032	277	529	280	1256	1214	199	440	106	1405	1284	296	814	287	997	7.49	35	1820	355	3426	ንሃዕ ንዳዳራ	0000	ר ר ר
10Fo	657 779 693 504																															
٦	0000																															
ᅩ	1111	7	ထောင	ထထ	ထတ	თ თ	6	ט כ	10	10	10	110	1.	11	11	12	7 7	12	13	13	13	- T	14	15	7	⊣,	<u> </u>				- ۱	1
ڃ	6 S C 6																							_	<u>~</u>	o 1	~ .	α α		-, ب		

_

																•																									
10s	94	45		-			20			29	18	59	33	25	35										36				21		21		80	67	16		17	20	34	3.7	19
10Fc	130	262	115	143	846	253	1690	1827	1699	1298	455	3149	2572	460	348	438	1383	2227	1750	79	931	1993	176	466	252	61	1.65	927	721	3367	355	1476	1579	4193	1284	25	962	1105	853	344	880
10Fo	94	236	0	116																										3154							σ.		1117	0	-
-	y y	ာဖ	9																																		9	9	9	9	9
ᅩ	٦,	i	_	7	7	7	7	~	7	7	7	7	7	7	2	٣	ო	٣	٣	٣	٣	r	m	. m	e	٣	4	4	4	4	4	4	7	4	4	4	4	4	2	S	2
ᇨ	20 1	- თ	11	-12	-10	8	9 1	-4	-2	0	7	4	9	æ	10	-11	6-	-7	15	-3	-	-	٣	5	7	9	-12	-10	-8	9 .	-4	7-)	?	4	9	80	10	-11	6-	-7
10s	23	31	36	41	17	29	44	17	16	17	21	109	57	44	27	23	15	14	28	32	19	38	75	21	41	28	18	22	26	25	77	65	670	36	11	11	29	145	23	49	41
10Fc	788	457	1727	1841	459	2060	3463	984	770	848	756	240	284	1404	1096	1349	1056	808	290	1977	066	492	143	1645	3297	1551	861	1844	2119	479	1540	248	3/3	17/2	1079	1195	1317	154	297	507	1219
10Fo	791																																								
1 1	r Cr																																								
يد	9	. ~	7	7	7	7	7	7	7	7	7	8	8	ဆ	8	8	æ	8	8	8	8	6	9	6	6	6	6	0	o.	o و	2 9	2 9	2 9	<u> </u>	0	0	0	01	0	Ξ	-
ч	10	16	۲-	-5	ų	-	1	က	2	7	σ	-10	8	9-	b -	-2	0	7	4	9	80	9	-7	-5	-3	-1	, 1	en ·	S)	· ·	p (0 <	יי ו	7 ())	7	4	9	ω	0 0	. L-
10s	24	35	44	34	61	56	28	30	29	21	117	25	29	24	40	7	26	പ	22	19	28	15	13	11	32	23	16	.26	;	97	7 0	97	10.	ם דום	35	13	14	91	23	36	39
	203																																			44	30	31	43	45	57
10Fc	_								∵	_																	1103												4	ň	5
10Fo	167	336	246	426	278	689	895	456	584	279	260	589	553	923	306	0	963	0	721	147	936	231	779	078	480	365	1071	127	0 0	089	907	000	ט מ	טינ טינ	215	544	670	76	457	310	709
1 1	4 1	4	4	4	4	4	4	4	4 1	4 1	4	4	4	4	4	Ŋ																_	7						ري ا	S	2
٠ بد	2 2	7	2	2	7	ლ -	<u>س</u>	<u>~</u>	<u></u>	E .	<u>.</u> 3	7	7	۲4	4	٠,	–			_	7	-	-	7	-	~	_	. .	N (7 C	7 C	1 c	1 0	4 (v (.V (7	0	7	7	m
ų	-4 1	0	2	4	. 0	. 5	€. 	7	-	<u>ო</u>	ro -	-4	7-	0	7	13	11	6	-7	2	61	-1		m	2	7	6		77) q	ט כ ו ו	> <	י ר ו	۱ د	> 0	7 '	4	9 (ω ,	10	11
																ı	1												I	1											1.
10s	198		17	14	91					1.5	,	-												37															20		16
10Fc	301	2657	1701	860	1664	2299	440	1100	099	1146	238	140	385	481	815	1598	587	338	118	1672	1155	302	10	844	393	122	943	36/1	CTT	1236	2793	5,7	745	000	000	3,00	101	1067	786	1210	426
0Fo	199 939	946	771	868	/31	447	496	045	657	084	280	144	111	787	810	019	597	390	75	589	068	267	0	915	616	0	075	979	777	780	765	2	708	200	7 0 0	אר ר אר ר	311	249	.072	354	483
1 1	4 4	4 2	4 1	4.	4.	4.	4.	-1	Φ.		<u>.</u>															4	4 1	مار مار	,	7 7	1 0	. 4	. 4	٠ -	, יי		•		4		
*	4 4	4	4	ማ •	7' '	4.	4.	4.	4	σ,	₫ (n ı	ດເ	ر ک	n ı	<u>ر</u>	Ωı	٠ ١	ഹ	n.	s i	ις.	2	9	9	9	9 (ه ر	ه د	ט כ	ی د	ی د	ی د	ט פ	י כ	- r	، -	~ r	- 1	- 1	_
æ	-12	-8	9	-4	7 (0 (7 .	4,4	9 ;	ω ,	10	11-	וע	- 1	r C	ا بىرى	⊣ ,	(m ı	n ι	۲ ,	o	11	-12	-10	8-	9-	1 4 (7 0	۰ د	1 4	ی د	α	10	1 -	7 7	ן אינ	- 1	n د	Ϋ́,	⊣
10s	35 25	35	20	46	220	77	30	36	53	7.7	31	7 7	, I	104	77	π ·	7 ,	134	;	23	31	٦ ;	56	20	43	56	40	ς. γ	20	4 9	·	۱	23	200) -	٦,	, ,	101	7 U	000	50
10Fc	1192 1679																																								
FO	1432																																								
10																								_																	
	m m																																								
~	3 10																																								
E	9 1	7-	,,,	۰ ر		• `	_ (~ (1	3	i	, -	' '	٠, (, .	. , ,	ų.	ĩ	ī	1	1	، ر	. ` '	7. '	- 1	Ϊ΄ '	1	1	-, (°	, u	- 4	2)	()	. <	ī	7 5	7 7 7	ם נוד נוד		í

0.8	32 24 34 34	26 70 27 52 34	22 22 22 22 22	50 50 50 50	33 33 33 23 23	25 25 25 25 25 25 25 25 25 25 25 21 21
0FC 1						932 366 1388 2254 708 734 748 304 440 11114 909
10Fo	51 32 66 137 18	87 255 141 19 3	108 210 210 135 135 171	242 6 100 47 17 34	480 1792 2044 0 621	134 134 134 134 104 104 111
7	00000	000000	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	. თ თ თ თ თ თ თ	
×	99999	00000		- 1 1 1 8		100 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0
£	-10 -8 -6 -4	07498	11.	1 3 7 -10 -8	0 4 2 0 2 4 8	
10s	33 32 24 19	19 32 27 60 29	20 22 22 31	33 35 32 32	46 30 37 37 96 1	158 30 . 22 64 32 16 19 19 11 14 25
10Fc	438 018 133 254 530	149 649 849 3330 783	363 363 965 704	535 53 53 863	466 967 170 530 273 37	543 237 1611 1018 74 1298 732 1932 1932 135 2220 960 960 925
0 1						
10F	48 100 113 125 59	1111 67 79 27 27 81	114 114 37 37 97	135 123 123 63	40 99 17 17 56 21 80	495 189 1581 965 64 1272 750 1681 146 2114 895 911
~	~~~~~	~~~~~~~			& & & & & & & & & & & & & & & & & & &	& & & & O O O O O O O O O O O O O
يد	00000	660000	170007	1	12 12 12 12 12 12 12 12 13 13 13 13	113
۳	3 1 1 1 1 2	2 - 8 9 - 1 - 1	1044064	5 3 3 4 4 6 6	04400445	0 7 2 3 3 3 7 5 7 6 7 7 8 7 7 8 7 8 7 8 7 8 7 8 7 8 7 8
		, , , ,	, ,			3 1 1 1 1
0s	25 17 24 36 38	36 13 13 40	23 24 29 17	19 19 16 23 .	78 42 42 25 28 23	34 20 22 22 21 33 33 11 11 11 11 11
ر. ري	. 0 0 H 2 H	пнань	1-04044	7 4 9 5 9 4	6674-198	5 1 8 0 2 8 2 2 2 5 4 2 0 2 5 5 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1
10Fc						2935 751 548 640 620 998 2282 522 1045 1972 1302
10Fo	644 889 759 912 659	993 962 277 721 721 587	993 459 007 007 178 305	325 325 331 778 503 0	202 305 1307 1557 1036 833 2984	929 707 707 519 689 689 689 689 689 737 737 906 972 247
7	2 1 1 8 8 8 8 × 1 1 8 × 1 1 8 8 × 1 1 8 8 × 1 1 8 × 1 1 8 × 1 1 1 1	88 11 8 11 11 11 11 11 11 11 11 11 11 11			8 8 1 8 1 8 1 8 2	88 1 2 1 8 1 8 8 8 8 8 8 8 8 8 8 8 8 8 8
<u>يد</u>		പലപപപ <i>റ</i>	1000000	100000		88884444444444
_	133	162760	1089470	24980	19673811	8 5 7 6 8 9 8 9 8 9 8 9 9 9 9 9 9 9 9 9 9 9 9
	- 5 - 1 - 1	โ	1 1 1 1 1	Ä	-11 -7 -7 -3 -1	201- 112- 124- 144- 144- 144- 144- 144- 14
Ø	12837	17.8983	0 9 2 9 8 7	850 0 - 1 0 -	အက္အဝတ္က က	24 19 33 33 20 20 24 14
10	2 2 2 4 4 2 1 2 2 4 4 2 2 1 1 2 2 1 1 2 2					-
10Fc	1958 28 72: 159:	16] 132. 166] 762	360; 2056; 56; 76; 76;	139 35 127 57 172	858 284 61 61 15 111 1111 418	733 2613 3221 1280 305 1767 691 469 212 362 656 758 1496
10Fo						743 3224 1240 298 1676 671 170 342 697 805
10	20 3 7 15	12	2001	12	288	1225 1166 1166 1155 1155
7						
24.			, o o o o o r	L L L L L L	L L L L B B	
æ	11 00 00 00 00 00 00 00 00 00 00 00 00 0	-109 -109 -16	0 2 2 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4 4	2 L L S L L L L L L L L L L L L L L L L	1 3 7 7 10 -10 -8	1 1 0 4 0 0 0 0 0 0 0 0
10s	21 18 15 19 25	62 25 29 29 21	103 30 44 32 23 23	21 23 14 48	22 32 32 41 41 1	2 2 3 3 3 3 5 7 1 1 1 8 8 8 7 7 8 8 8 8 9 7 9 8 8 9 9 9 9 9 9
10Fc	1188 1607 1045 666 590	213 792 449 1042 410 826	108 693 391 1540 808 958	810 892 107 429 54	968 333 160 613 262 84	988 1111 474 364 821 1037 770 770 25 867 524 524 1210
.0Fo 1			104 689 400 .473 943			
		_			_	958 1010 1010 1010 1010 1010 1030 1130
-						0000000000000000
×	000000	100000000000000000000000000000000000000	наннан			
ч		8 - 1 - 6 2 - 1 - 2 2 - 2 - 2	6 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 8 9 1 6	20 0 4 5 6 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	1 1 2 2 2 2 1 1 1 2 2 2 2 1 1 1 1 2 2 2 1

																													٠														
10s	,	40	2,5	23	20	; -	32	2 00	7 1	30	7	25	27	20	37	16	34	29	24	30	39	-	39	33	52	37	64	32	30	151	56	24	۲,	⊣ (70		52	80	22	56	48	34	47
0Fc	6.0	628	991	632	1032	144	869	593	517	709	107	663	367	1036	846	219	707	2036	1438	375	186	164	846	966	311	571	291	1443	9201	125	784	629	200	177	340	757	200	313	973	229	825	325	456
0Fo 1																																									963		
1 1	,		2 2	13 1								51																			<u>.</u>	ກ :	2.5	2 5	2 5	or 5	7 7	7 2	.4	14	7	14	7
ᅶ	ų	9 9	9 49	9	9	9	7	_	-				~	7	8	ω	80	89	ω	80	8	ου	9	6	6	6	6	10	10	10	0.1	07.	 	7 .	+ 0	- ·	, .	_ _	0	0	0	0	0
E	1	- 2	0	5	4	9	6-	-7	ي ا	1	1	-	m	5	8-	9 .	-4	-2	0	7	4	7-	-5	-3	-1	-	c	9-	-4	-2	> (7 4	י נ	٠. د	T (0.7	0 .	9	-4	-2	0	7	4
10s	,	25	63	26	47	23	22	7	2.7	35	49	56	23	22	27	34	19	29	27	22	31	21	42	79	37	62	24	39	٦	37	٦ (ر د د د د	55	72	7 C	0 0	, ,	200	27	52	25	16	12
0Fc																																									836		
.0Fo 1																																									783		
1	12	12	12	12]	12	12	12	12	12	12	12	12	12]	12	12	12	12	12	12	12	12	12	12	12	12	12	12	12	175	7.7	12	12	1 0	12 7	1 5	7	1 .	77	۲۲:	13	13		13
. بد	G	9	9	9	9	9	9	7	7	7	7	7	7	7	7	æ	80	89	ω	œ	œ	œ	6	6	9	σ	σ,	٥ ا	10	2.5	2 5	2 2	-	; [; ;	1 -	1,5	71	٦ ,	∸.		-4 ,	7
ŗ	9-	-4	-2	0	2	4	9	6-	-7	-5	i,	-1	~	m	S	-8	9-	-4	-2	0	2	4	-7	7	۳ ا	Ţ	, , ,	, ن	9 ,	i i	7 0	<i>ر</i>	ו) ר ו) (- 1	1 6	7 -	11-	ו זע	7-	n c	2
10s	28	99	101	49	35	28	34	25	31	30	87	69	25	39	24	48	35	70	23	28	47	45	39	47	٦ ;	34	31	- :	7.5	ا ا	۲. ۲.	36	62	101	31	20	, 6	2 6	יים ת	35	22	67	٦ ۲
10Fc	946	316	81	1355	2272	166	419	1192	1556	1303	208	48	1301	404	828	521	970	518	351	1243	1290	1472	782	795	126	909	1035	24	21/9	223	1022	2415	267	270	778	870	0.56	2 5 5 5	44.000	ימת ימת	169T	100	400
.0Fo				_		_	_				_	_		_	٠.				_		_	_			_			_		_											1593		
-	11	11	11	11	11	11	11	11	Ξ.	11	11	Ĺ	11	11	11	11]	7	Ξ,	Ī	11		11	Ξ:	12	77	77	7 7	77.	12	12	12 2	12	12	12	12	12 1	1 5	10 6	7 7 7 7	1 21	12	7
ㅗ	9	6	9	9	0	σ	6	9	10	10	10	10	10	10	10	11	17	11	Ξ:	7		7.7	12	7.7	0 0	> 0	> c	٥	o c	· C	0	0	0	0	Н	-	~	· -		⊣ ⊢	٦ ,-	٠.	4
æ	6-	-7	-5	-3	-1	Τ.	en.	5	8	9 -	-4	-2	0	7	4	-7	7,			_	m ·	b	2	٠,	-12	-10	χ (1	P	ן ו	7 0	2	1 4	9	80	-11	6-	7-	ی ا	יי) ל !	ا ا	- -	÷ (*	,
10s																																									41		
10Fc	736	909	166	447	940	295	206	439	692	235	1086	243	1484	964	2289	504	920	, 68	/611	108	7.00	171	215	332	717	700	100	750	6.5.1	202	680	196	1554	86	1375	1612	1938	9	573	92.5	195	1071	• •
10Fo	795	630	153	441	902	301	188	459	646	316	1071	261	1363	. 881	8802	483	4.4	700	9911	0 0	243	080	239	400	120	010	107	286	07	192	680	198	1489	92	1282	1559	2016	0	590	927	41	1018	} }
7	10	10	10	10	10	10	10	10	10	11	11	11	7.7	11	- ;	- ;	11	4 -	- -	Ϊ,	T .	7 ,	7 -	 	7 -	4 c	11	; =	; -	-	11	11	11	11	11	11	11	11		; ;	11	11	j I
×	11	11	12	12	12	12	12	13	13	1		Η,	-		→ .	٦,		, -	⊣ (V (7 (7 (7 (7 (7 0	4 0	۱ ر	1 0	1 ~	ı m	e	æ	m	3	m	c	က	m	, m	۷ ۲	4	4	
ב	1	m '	9-	-4	-2	0 (2	ŗ,		-11	6			1	7 -		יטרי	יו כ	- 6	77	07-	0 (0 5	7 (7 C	,	1 4	. (c	œ	-11	6-1	1-	-5	-3	-1	-	m	5	7	-10	φ • •	1	
10s																																									73		
10Fc	2465	3839	1231	356	1442	1430	204	576	849	284	94	490	341	1206	1074	1014	000	37.	306	1170	6111	2007	700	2350	2916	422	345	630	782	87	51	828	1132	873	80	374	75	780	1052	335	173	1446	
10Fo	2250	3774	1175	386	1397	1.408	9/1	. 585	88.5	254	۰.	469	27.7	114	1001	199	757	נטנ	767	700	2457	- 6 - 6 - 6 - 6 - 6 - 6 - 6 - 6 - 6 - 6	607	2268	2884	434	391	684	810	110	61	752	1038	822	0	395	117	802	1033	319	186	1371	
7	10	10	10	10	707	0,7	0 7	10	2	10	10	10	2 .	7 0	2 6	7 0	2 5	2 -	2 -	9 6	2 -	2,5	2 5	2 -	10	2 -	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	10	
×	7	2																		•	r <	۰ <	7 9	• 4	4	. 4	4	4	2	2	2	Ġ	S	S	വ	S	'n	2	9	9	9	9	
ч	-4	7 9	0	N •	4, 4	ه د	χ, α	77-	וע	7-	<u>۱</u>		1	٦ ٣	ייר	י נ	٠ ٥	-12	101-	2 4 1	י ו	1	r C	1 0	2	4	9	89	-11	6-	1-1	-5	ر 1	-1	7	د	S	7	-10	-8	9-	-4	

10s	ç	7 6	7 4	-	104	3	, -	4 1	7 6	י ע	ט כ	3.5	י מי	50	7 (39	39	36	37	23	2 0	20	5 6	9	41	34	42	38	36	34	59	25	100	43	9 C	۳,	49	89	3	53	98	4. د ر	4°	79
10Fc	100	770	321	181	500	845	133	758	086	2021	999	1.4	597	438	462	613	603	983	1107	759	200	345	595	929	712	1019	438	617	773	691	731	888	196	638	179	36	408	1154	419	572	257	514	6/5	yaa yaa
10Fo																															918													
1 1	,		_		7		7																																	3 0 (ם ת	אָרָ ק	7
×	ď			9	7	7		. &	0 00		, -	0		0		-	-	-	-		2	2	2	. ~	2	2	ω,	3	3	3	3 18	4	4.	7 ,	.	ח.	ດ ເ	η. Ο (ָ י	9,				7
ď	١	1 0	-2	0	5-	٦ (1	-4		1 α	י ו	-4	- 2	0		1-	-5	3	-	-	8	9	-4	-2	0	7	-7	-5	-3	-1	7	9	1 4	7 0	-	ດ	י ר	⊣ •	ਰਾ (ਹ	7 7	· ·	0 0	ו ו	4
10s	0	27	34	70	47	34	24	41	·	31	125	152	41	27	92	33	9	97	-	29	66	43	Н	35	38	-	48	41	44	35	42	70	ν . Σ. (70	0 7 0	27. כר	ς.	۲ ,	70	۲. c	000	0 K	י ע מ	2
10FC	111	1497	705	154	366	917	979	328	95	818	27	201	756	1120	174	837	294	446	86	648	335	611	55	460	618	122	337	882	468	891	492	1041	000	202	143	141	130	100	7 (300	700	907	195)
0Fo :																															634													
1 1	16.1	16.1	16																												17													
*	4	4	2	ß	ß	'n	S	ω.	ß	9	9	9	9	9	9	7	۲.	7	_	۲.	ω	8	ω	8	თ	6	, ,	-		-			- ·	4 0	1 0	1 0	1 0	4 6	1 0) r	, ~	, .	, m	,
£	0	4	6-	-7	ı S	m 1	-1	1	c	8-	9	-4	-2	0	7	-7	5	-3	7		9-	-4	-2	0	-3	-1	6-	-1	-5	ا ع	Τ,	⊣ c	nα	9 4	1	- 6	1 C	, ,	1 0	, ,	ی ۔ ا	1	1	ı
10s	36	98	35	55	-	21	29	29	23	64	22	19	96	39	1	25	21	163	29	57	44	113	93	9 6	22	69	52	7.1	42	99	ю . Н	000	ر م م	52	ı -	46	3.6		- 6	200	4 4	41	30	ı
10FC																															1070													
.0Fo]																															1111													
1	1.5	15	15	15	15	15	15 1	15 1	15	15	15 1	15 1	15	15	15	15	15	15	15 1	15 1	15	15	16	. 91	16 1	16. 2	16	, , i	167	7 07	 	7 7	1 9	16	9	16	9	9	9	9	16 1	16 1	16	
74	9	9	9	9	9	9	7	7	7	7	7	7	ထ	æ	æ	ထ	œ	თ	9	0	6	10	0	0	0	0	0	0	-	۰ د		٠.	-	-	-	-	~	۱ ۸	٥.	2	7	7	7	
ď	91	-4	-2	0	7	4	-1	-5	-3	-	1	æ	9-	-4	2	0	7	-5	٦.	7	7	-2	-10	-8	9-	-4	-2	0 (7 •	ਬਾ (אינ	۱ ۱	1	- 1	_	ľ	-10	000	9	-4	-2	0	7	
10s	-	19	7.1	33	~	54	46	23	103	45	120	31	37	45	156	39		21	26	17	29	33	83	32	28	84	27	c	200	0 0	12	. 4.4	45	31	26	89	27	37	20	35	61	145	45	
10Fc	167	907	480	892	.86	331	583	579	386.	624	138	865	439	541	81	655	134	860	276	1094	937	1167	202	916	732	102	411	25	507	707	ر د د د	361	259	682	1604	73	410	1130	1647	414	644	22	425	
10Fo	0	936	478	784	0	280	602	211	301	522	120	871	420	563	173	579	0	967	267	1096	968	1142	166	915	780	82	436	ט	0/7	27.0	; C	363	256	695	1615	69	350	1171	1706	446	577	146	438	
-	14	14	14	14	14	14	14	14	14	14	14	14	14	14	14	7.	14	15	12	15	12	12	1.5	15	15	15	٠ ر	12	0 4	י ב	7 .	15	15	15	15	15	15	15	15	15	15	15	15	
*	7	7	œ	8	œ	ထ	æ		-			6		Ä,	H	Ä,	4			Α,	<u> </u>	٠,		Н.	7	7	7 (ν ς	4 C	4 C	1 0	1 2	c	٣	m	٣	٣	e	æ	m	4	4	4	
æ	7	e	-8	9 -	-4	-2	0	2	-7	1.5	-3	7	1	9-	-4	7.	•	ן טנ	- 1	-5		[,	٦ ،	m i	ດຸ	-10	i S	9 7	, (1 C	0	4	6-	-7	-5	-3	-1	1	3	5	-10	-8	9-	
10s	35	71	-	37	20	21	40	23	24	29	24	20	39	38	43	121	2.5	24	C 7	٠,	٦ ;) C	17	29	48	η·	יר עינ	110	23	7 4	53	32	38	33	25	24	51	34	24	91	46	22	64	
10Fc	1053	220	37	313	597	1006	826	785	874	620	1481	790	108	2161	6/67	887	0 2 2	067	4	114	900	223	278	711	751	919	1167	186	1135	1895	259	480	1057	1234	594	708	357	602	121	72	1038	762	202	
10Fo	1101	192	0	285	617	1028	840	764	926	628	1448	771	40	2133	2363	121	770	00/	400	٥	0 0	155	307	707	167	000	7 6 6	7	132	730	212	489	174	265	611	702	340	632	24	92	983	810	159	
-	14]	14	4	4	4	4	4	4.	4	4	4	σ,	4.	4	4.	a	r <	Jr <	, ·	. -	7 4	ਰਾ <	J .	.	₹ <	. .	. <	. 4	. 4	. 4	14	4	4	4	4	4	4	4	4	4	4	4	4	
×	0	-										~ (4	4	4	4	4	4	4	ß	2	2	S	ഗ	ر د د	S	
ч		-11						~ (m i			æ v				c د	1 -	# U	9 -	→ C	ז ת ו	<u>۔</u> ا) (י ני	 I	⊣ ` ດ	า ช	, -	α α	9	-4	-2	0	7	4	9	6 -	۲.	-5	.3	-1	- 1	~	,

	131	5	2	8	,	31	36) (971	ď	מ	80	, (20	8 U	9				
								, (
													•	•						
	7	10	1	2 19		7	3 19		עיי	3 19		4 19	0	7	1 20	1				
								,												
22	7	23)	30	101	T 7 T	34	8 7	r	65	, (99	7	7	112	0	9	٠ ر	9.	~
617	10	1008		827	201	707	597	000	000	681	1 (268	697	5	72	000	780	864		322
								861												
~	,	ς.	, ,	4	4	• •	4	4 17	,	7		J'	G.) (ر ر	ď	7	3) (
_		m		ρ	9	٠ ،	14	-2	1	0	c	7	-7	٠ (ť,	۲	י	ì	۰,	-
								99												
								156												
126		17	000	700	599		717	100		353	89	3	332		0101	1695		133	000	200
								3.16												
2		_ກ	٠	1	m	٠	າ	د		ກ	~)	4	*	.	4		4		.
4		1	-	ī	15	,	ו	-	-	_	~	,	8		i	-4	•	-2	_	•
2.4		-	~	7	35	-	7	30	0	40	_	•	24	20	, ,	99		20	32	1
								978												
								806												
7	1	., 	4 15		4 15	1 0	-	5 15		·	5 15		2	7)	5 15		C	9	,
								l D												
								73												
7	340	ה ה	128	,	1122	1740	-	9	909		563	000	976	44		1045	7.77	5	499	
,	œ	•		,	٥	9	, ,	6 14	œ	, ,	9	,	-	7	,	_	7		_	